arXiv:1603.06842v1 [nucl-th] 22 Mar 2016

Role of spd- Continuum Components in the Halo Nucleus ^{6}He

Jagjit Singh[1,2,a)]

[1]*Dipartimento di Fisica e Astronomia "G.Galilei", via Marzolo 8,I-35131 Padova, Italy*
[2]*INFN-Sezione di Padova, via Marzolo 8, I-35131 Padova, Italy.*

[a)]Corresponding author: jsingh@pd.infn.it

Abstract. Role of different continuum components in the weakly bound nucleus ^{6}He is studied by coupling unbound spd-waves of ^{5}He by using simple pairing contact-delta interaction. The results show that ^{6}He ground state 0^+ displays collective nature by taking contribution from five different oscillating continuum states that sum up to give an exponentially decaying bound wavefunction emerging from five different possible configurations i.e. $(s_{1/2})^2$, $(p_{1/2})^2$, $(p_{3/2})^2$, $(d_{3/2})^2$ and $(d_{5/2})^2$. The ground state properties of ^{6}He has been calculated.

INTRODUCTION

The nucleus ^{6}He is a typical "neutron skin or halo"[1] with a very low neutron pair separation energy (0.975 MeV) [2]. Motivated by the recent experimental measurements at GANIL [3] on continuum resonances in ^{6}He, recently we have developed a simple theoretical model [4] to study the weakly bound ground state and low-lying continuum states of ^{6}He by coupling two unbound p-waves of ^{5}He. In the present study we have extended the model space with inclusion of sd- continuum waves of ^{5}He. The large basis set of these spd- continuum wavefunctions are used to construct the two-particle ^{6}He ground state (g.s.) 0^+ emerging from five different possible configurations i.e. $(s_{1/2})^2$, $(p_{1/2})^2$, $(p_{3/2})^2$, $(d_{3/2})^2$ and $(d_{5/2})^2$. The simple pairing contact-delta interaction is used and pairing strength is adjusted to reproduce the bound ground state of ^{6}He. The extension of model space is a computationally challenging problem. The main aim is to show how an extension of theoretical concepts related to residual interactions, namely a contact delta pairing interaction, naturally explain the stable character of the bound states of Borromean nuclei, such as ^{6}He and simultaneously account for some of the resonant structures seen in the low-lying energy continuum.

SPECTRUM OF ^{5}He

The unbound nucleus ^{5}He can be described as an inert ^{4}He core with an unbound neutron moving in p doublet or d doublet or s singlet in simple independent-particle shell model picture. These doublets are further splitted by spin-orbit interaction. Experimentally only the $p_{3/2}$ and $p_{1/2}$ resonances are confirmed at 0.789 and 1.27 MeV above the neutron separation threshold and their widths are quoted as 0.648 MeV and 5.57 MeV respectively [5]. Theoretically in order to extend the model space we have also included the $sd-$shell in picture. The continuum monopole ($\ell = 0$), dipole ($\ell = 1$) and quadrupole ($\ell = 2$) scattering single particle states ($E_C > 0, k > 0$) of ^{5}He (see Fig-(1)) are generated with Woods-Saxon (WS) potential given by

$$V_{WS} = \left[V_0 + V_{ls} r_0^2 (\vec{l}.\vec{s}) \frac{1}{r}\frac{d}{dr}\right]\left[1 + exp\left(\frac{r_0 - r}{a}\right)\right]^{-1} \quad (1)$$

where $r = r_0 A^{\frac{1}{3}}$. For ^{5}He the parameter set used is WS potential depth $V_0 = -42.6$ MeV, $r_0 = 1.2$ fm, $a = 0.9$ fm and spin-orbit coefficient $V_{ls} = 8.5$ MeV.



FIGURE 1: (Color online) ^{5}He continuum waves as a function of radial variable for continuum energies 1, 5 and 10 MeV respectively

MODEL AND RESULTS

The simple model with two non-interacting particles in the above single-particle levels of ^{5}He produces different parity states (see TABLE-1) when two neutrons are placed in five different $s_{1/2}$, $p_{1/2}$, $p_{3/2}$, $d_{3/2}$ and $d_{5/2}$ unbound orbits. Namely five configurations $(s_{1/2})^2$, $(p_{1/2})^2$, $(p_{3/2})^2$, $(d_{3/2})^2$ and $(d_{5/2})^2$ couple to $J = 0^+$.

TABLE 1: Possible configurations of ^{6}He arising from two neutrons in s-, p- and d-orbitals

	$s_{1/2}$	$p_{1/2}$	$p_{3/2}$	$d_{3/2}$	$d_{5/2}$
$s_{1/2}$	0^+	$0^-, 1^-$	$1^-, 2^-$	$1^+, 2^+$	$2^+, 3^+$
$p_{1/2}$		0^+	$1^+, 2^+$	$1^-, 2^-$	$2^-, 3^-$
$p_{3/2}$			$0^+, 2^+$	$0^-, 1^-, 2^-, 3^-$	$1^-, 2^-, 3^-, 4^-$
$d_{3/2}$				$0^+, 2^+$	$1^+, 2^+, 3^+, 4^+$
$d_{5/2}$					$0^+, 2^+, 4^+$

The two-particle wave functions are constructed by tensor coupling of two continuum states of ^{5}He. The five states discussed above are not discrete, but rather depend on the energies of the continuum orbitals. Each single particle continuum wavefunction is given by

$$\phi_{\ell,j,m}(\vec{r}, E_C) = \phi_{\ell,j}(r, E_C)[Y_{\ell m_\ell}(\Omega) \times \chi_{1/2,m_s}]^{(j)}_m \tag{2}$$

The combined tensor product of these two is given by

$$\psi_{JM}(\vec{r}_1, \vec{r}_2) = [\phi_{\ell_1,j_1,m_1}(\vec{r}_1, E_{C1}) \times \phi_{\ell_2,j_2,m_2}(\vec{r}_2, E_{C2})]^{(J)}_M \tag{3}$$

For simplicity an attractive pairing contact delta interaction $(-g\delta(\vec{r}_1 - \vec{r}_2))$ is used, because we can reach the goal with only one parameter adjustment. The continuum single-particle wavefunctions are calculated with energies from 0.0



FIGURE 2: (Color online) Eigenspectrum of the interacting two-particle case for $J = 0$ for increasing basis dimensions, N. The coefficient of the $\delta-$contact matrix, G, has been adjusted each time to reproduce the g.s. energy (right). The actual strength of the pairing interaction, g, is obtained by correcting with the energy spacing ΔE and it is practically a constant.

to 10.0 MeV and normalized to a delta for the spd-states of ^{5}He on a radial grid which varies from 0.1 fm to 100.0 fm with the potential discussed above. The two particle states are formed using mid-point method with an energy spacing of 2.0, 1.0, 0.5, 0.2 and 0.1 MeV corresponding to block basis dimensions of N =5, 10, 20, 50 and 100 respectively and the matrix elements of the pairing interaction are calculated. In Fig-2, the eigenspectrum for $J = 0^+$ case is presented and from figure it is clear that with increase in basis dimensions the superflous bound states moves into the continuum. The coefficient of the $\delta-$contact matrix (G) is adjusted to reproduce the correct ground state energy. The actual pairing interaction g is obtained by correcting with a factor which depends on the aforementioned spacing between energy states. The correction factor is practically a constant quantity, except for the smallest basis. The biggest adopted basis size gives a fairly dense continuum in the region of interest. The radial part of the $S = 0$ ground state wavefunction obtained from the diagonalization in the largest basis is presented in the left part of Fig-3. It shows a certain degree of collectivity, taking contributions of comparable magnitude from several basis states. The surface plot shows the exponential behavior typical of a bound state, despite being the sum of many products of oscillating wavefunctions. One can see from the right part of the Fig.-3 that the square of the amplitudes of the $(p_{3/2})^2$ components are dominant summing up to 89.7%. The comparison among present results and earlier studies by T.Myo [6] and Hagino [7] is presented in TABLE-2 and from this table it is clear that present results are in good agreement with earlier calculations, which further strengthen present results. In TABLE-3, calculated ground state properties are compared with earlier studies[6, 7], where R_m is the matter radius,

$$\langle r_{NN}\rangle = \langle \psi_{gs}(\vec{r}_1,\vec{r}_2)|(\vec{r}_1-\vec{r}_2)^2|\psi_{gs}(\vec{r}_1,\vec{r}_2)\rangle \tag{4}$$

is the mean square distance between the valence neutrons, and

$$\langle r_{c-NN}\rangle = \langle \psi_{gs}(\vec{r}_1,\vec{r}_2)|(\vec{r}_1+\vec{r}_2)^2/4|\psi_{gs}(\vec{r}_1,\vec{r}_2)\rangle \tag{5}$$

is the mean square distance of their centre of mass with respect to the core.



FIGURE 3: (Color online) Ground state wavefunction ($S = 0$) for N =100 as a function of the coordinates of the two neutrons and corresponding contour plot (Left part). Decomposition of the g.s. into the J=0 basis (Right part) as a function of an arbitrary basis state label $|i>$: the basis is divided into five blocks, 10^4 $[s_{1/2}\times s_{1/2}]^{(0)}$, 10^4 $[p_{1/2}\times p_{1/2}]^{(0)}$, 10^4 $[p_{3/2}\times p_{3/2}]^{(0)}$, 10^4 $[d_{3/2}\times d_{3/2}]^{(0)}$ and 10^4 $[d_{5/2}\times d_{5/2}]^{(0)}$ components.. The ordering in each block is established by the sequential energies of each pair of continuum s.p. states, i.e. (E_{C_1}, E_{C_2}) = (0.1, 0.1), (0.1, 0.2), . . . ,(0.1, 10.0), (0.2, 0.1), (0.2, 0.2), . . . (10.0, 10.0).

Config.	Present	T.Myo[6]	Hagino[7]
$(2s_{1/2})^2$	0.008	0.009	–
$(1p_{1/2})^2$	0.080	0.043	–
$(1p_{3/2})^2$	0.897	0.917	0.830
$(1d_{3/2})^2$	0.005	0.007	–
$(1d_{5/2})^2$	0.009	0.024	–

TABLE 2: Components of the ground state (0^+_1) of ^{6}He

	Present	T.Myo[6]	Hagino[7]
R_m	2.37674	2.37	...
r^2_{NN}	28.8404	23.2324	21.3
r^2_{c-2N}	7.21011	9.9225	13.2

TABLE 3: Radial properties of the ground state of ^{6}He in units of fm

CONCLUSIONS

In the present study I present emergence of bound halo ground state of ^{6}He from the coupling of five unbound spd-waves in the continuum of ^{5}He, due to presence of pairing interaction. Contribution of different configurations has been presented. Radial properties of ground state of ^{6}He are also presented and compared with other calculations.

ACKNOWLEDGMENTS

The useful discussions with L.Fortunato, A.Vitturi, R.Chatterjee and Sukhjeet Singh and financial assistance from Fondazione Cassa di Risparmio di Padova e Rovigo (CARIPARO) is gratefully acknowledged.

REFERENCES

[1] I. Tanihata *et al.*, Physics Letters B **160**, 380 – 384 (1985).
[2] F. Ajzenberg-Selove, Nucl. Phys. A **490**, 1 – 225 (1988).
[3] X.Mougeot *et al.*, Physics Letters B **718**, 441 – 446 (2012).
[4] L.Fortunato, R.Chatterjee, J. Singh, and A.Vitturi, Phys. Rev. C **90**, 064301–1, 6 (2014).
[5] TUNL, Nuclear data evaluation, `http://www.tunl.duke.edu/NuclData/General_Tables/5he.shtml` (2009).
[6] T. Myo *et al.*, Progress in Particle and Nuclear Physics **79**, 1–56 (2014).
[7] K. Hagino and H. Sagawa, Phys. Rev. C **72**, 044321–1, 6 (2005).